

# ON SOME OVIPAROUS FILARIAL NEMATODES MAINLY FROM AUSTRALIAN BIRDS<sup>(1)</sup>

by

ODILE BAIN\* and PATRICIA M. MAWSON\*\*

\*Laboratoire de Zoologie-Vers, associé au CNRS, Muséum national d' Histoire Naturelle, 43 rue Cuvier, F 75231 Paris Cedex 05

\*\*Department of Zoology, University of Adelaide, G.P.O. Box 498, Adelaide, South Australia 5001

## ABSTRACT

BAIN, O., and MAWSON, P. M. 1981. Some oviparous filarial nematodes mainly from Australian birds. *Rec. S. Aust. Mus.* 18(13): 265-284.

Three groups of filarial nematodes are represented in this collection.

### 1. Aproctinae.

Four species are described: *Pseudaprocta copemani* n.sp. from *Petroica multicolor*; *Aprocta boulengeri* n.sp. from *Strepera graculina*; *Aprocta bakeri* n.sp. from *Corvus orru*; *Lissonema* sp. from *Ninox novaeseelandiae*.

*Pseudaprocta* is regarded as being closely linked with the Scuratoidea.

The species assigned to *Aprocta* are heterogeneous, falling into two distinct morphological groups, one of which corresponds with the genus *Lissonema*, which is therefore reinstated. *Aprocta* spp. have a short ovejector, small caudal papillae in males, spicules generally subequal with fine distal extremities, no deirids, small eggs (less than 55 µm long), short larvae (around 200 µm long) with short tail bearing terminal spines. *Lissonema* spp. have a long ovejector, at least one large caudal papilla in male, spicules usually equal, with large blunt distal extremities, deirids, large eggs (more than 65 µm long), long larvae (more than 400 µm long) with long unarmed tail. The genus *Squamofilaria* is regarded as a synonym of *Lissonema* and its species are shown to belong either to *Lissonema* or *Aprocta*. *Lissonema* contains 20 spp. (19 n. comb.) and *Aprocta* 17 spp. (2 n. comb). The host ranges of these two genera are completely different.

### 2. Dicheilonematinae.

*Serratospiculum guttatum* (Schneider, 1866) from *Falco longipennis* and *F. peregrinus*, and *S. tendo* (Nitzsch, 1819) from *F. peregrinus* are redescribed. *S. seurati* n.sp. is described.

*Serratospiculum*, parasites of Falconidae, contains a total of seven species. These species fall within two groups depending on the spicule length. (1) Those

with short spicules are separated further, in that *S. turkestanicum* Skrjabin, 1916, *S. guttatum*, *S. kwangsiensis* Hsü, 1963, possess specific cuticular ornamentations in the female; while *S. chungii* Hoeppli et Hsü, 1929, *S. seurati* n.sp. (= *F. attenuata* Rud., 1819 sensu Seurat, 1915), *S. congolensis* Vuylstecke 1957, are distinguished by the morphology of the epaulettes and the right spicule; (2) those with long spicules: *S. tendo* (= ? *S. thoracis* Tubangui, 1934; = ? *S. lii* Ezzat et Tadros, 1958).

### 3. Diplotriaeninae.

Three species are described: *D. spratti* n.sp. from *Oreoica gutturalis*, *D. beveridgei* n.sp. from *Corvus orru*, and *D. smithi* n.sp. from *Acanthogenys rufogularis*. Additional morphology is given for *D. delta* Johnston and Mawson, 1940 and *D. falconis* (Connal, 1912).

## DEPOSITION OF TYPE MATERIAL

Holotypes and allotypes have been deposited in the South Australian Museum, Adelaide; (SAM in text); other material has been deposited as noted, in the Muséum National d' Histoire Naturelle, Paris (MNHN) and in the Australian Helminthological Collection (AHC) at present housed in the South Australian Museum.

In the species described here, the female has been designated as holotype, and the male as allotype. This is because the family, genera and species of filarial worms are more often identified by the female than by the male.

## DESCRIPTION OF MATERIAL

### I. Aproctoidea—Aproctinae

#### Genus *Pseudaprocta* Schikhobalova

The morphological characters of the genus *Pseudaprocta* are closer to those of Scuratoidea Ascaridida than of Spirurida (Chabaud, 1974): mouth large and triangular, six external labial papillae, situated on the same circle as the cephalic papillae, oesophagus short and very thick, without

<sup>(1)</sup> This study was carried out while working under a grant (to O.B.) from the World Health Organisation.

differentiation into two parts, spicules equal and simple, and one pair of caudal papillae situated laterally (8th pair).

It seems that *Pseudaprocta*, and possibly the Aprocinae, evolved from the Seuratoidea.

***Pseudaprocta copemani* n.sp.**

(Figs. 1, 2.)

**Material examined**

From *Petroica multicolor* Gmelin (Muscicapidae) from Maggs Mt, Tasmania. Holotype ♀, allotype ♂ (SAM V2096, V2097), 1 ♀ and 1 ♂ (MNHN, Paris, No. 201 HD); 2 ♀ and 3 ♂ (AHC 6277).

From *Pachycephala pectoralis* Latham (Muscicapidae), from Heatherleigh, S. Aust.: 3 pieces of ♀ (2 anterior, 1 posterior), 1 ♂, 1 posterior end ♂. (AHC 6282).

**Description**

Morphology is shown in Figures 1 and 2. The specimens, except in one case, are surrounded with a

thin membrane detached from the rest of the cuticle and forming a vesicle around the body, beginning at the level of the oesophagus and ending at the level of the tail.

In the female, the tail ends in a small rounded knob (Fig. 1, I). In the male, the precloacal unpaired papilla is particularly big (Fig. 2, B and C). The spicule sheaths are thickened near the cloaca; these thickenings apparently assume the role of a gubernaculum (Fig. 2, E).

**Measurements**

Holotype ♀: body length 16.5 mm, width 500 µm; nerve ring, deirids and vulva at 160 µm, 320 µm and 370 µm from head; length of buccal capsule 25 µm; length of oesophagus 585 µm; length of tail 320 µm.

Allotype ♂: length of body 8.9 mm; width 310 µm; nerve ring 160 µm from head; length of oesophagus 520 µm; length of left spicule, right spicule and tail respectively 640 µm, 600 µm, 190 µm.




Figure 1. *Pseudaprocta copemani* n.sp., female. A, anterior end, ventral view; B, C, & D, head in apical, dorsal and ventral views; E, larva; F & G, head of larva, from hook side and opposite side; H & I, tail in lateral and ventral views; J, tip of tail; K, transverse section of body. A, H, I, J, K, to scale 200 µm; B, C, D, to scale 100 µm; E, F, G, to scale 10 µm.




Figure 2. *Pseudaprocta copemani* n.sp., male. A, anterior end, lateral view; B & C, caudal region, lateral and ventral views; D, region of deirid and excretory pore, lateral view; E, distal extremities of spicules and gubernaculum, lateral view. A, B, C, to scale 200  $\mu$ m; D, E, to scale 100  $\mu$ m.

Paratype ♀: body length 18 mm; width 520  $\mu$ m; nerve ring, deirids, excretory pore and vulva at 152  $\mu$ m, 320  $\mu$ m, 270  $\mu$ m and 435  $\mu$ m from head; buccal capsule 25  $\mu$ m long; egg 52  $\mu$ m  $\times$  42  $\mu$ m; larva 480  $\mu$ m  $\times$  8  $\mu$ m.

Another ♂ from AHC6277: body length 9 mm long, width 400  $\mu$ m; nerve ring, deirids and excretory pore 160  $\mu$ m, 290  $\mu$ m and 230  $\mu$ m from head; buccal capsule 22  $\mu$ m long; oesophagus 440  $\mu$ m long; length of left spicule, right spicule and tail respectively 550, 575 and 200  $\mu$ m.

#### Discussion

This material is distinguished by its longer spicules from *P. gubernaculum* Schikhobalova, 1930 (Caucasia); *P. decorata* Li, 1933 (China); *P. buckleyi* (Singh, 1949) and *P. leiperi* (Rasheed, 1960) (India), and *P. ungriai* Chabaud *et al.*, 1964 (Madagascar), in which the spicules are about 400  $\mu$ m long. It differs from *P. sichotealinensis* Oshmarin & Belouss, 1951 from Eastern USSR (Sonin, 1966) and *P. skrjabini* (Ali, 1956) from India, by the rather greater spacing, longitudinally, of the papillae near the cloaca. *P. myzanthae* Johnston & Mawson, 1940, from

*Myzantha flavigula* (Meliphagidae) from South Australia, is known only by the female, the measurements of which are very similar to those of the present specimens with the exception of the body length which is greater. It seems better to keep the two species separate, in view of the difference in hosts, and of the fact that Johnston & Mawson show a slightly different arrangement of the cephalic spines (Johnston & Mawson, 1940, fig. 14). We therefore propose it as a new species, *P. copemani*, in honour of our Australian colleague Bruce Copeman.

#### The Genera *Aprocta* Linstow, *Lissonema* Linstow, and *Squamifilaria* Schmerling

The genus *Aprocta* according to Anderson and Bain (1976), is large and heterogenous. Some authors (Boulenger, 1928; Sandground, 1933 etc . . .) had previously proposed to recognize two genera, *Aprocta* Linstow, 1883 and *Lissonema* Linstow, 1903, separated on the form of the head (flat or with two lateral elevations), the oesophagus (divided or not), the caudal papillae of the male (few and small or large and numerous). As pointed out by

Anderson and Chabaud (1958) the two first of these characters are of doubtful use, because there are many species in which the division is not clear. However, since their work, further descriptions and redescriptions have been made (Ezzat and Tadros, 1958; Rasheed, 1960; Diaz-Ungria, 1963; Chabaud *et al.*, 1964; Schmidt and Kuntz, 1970; Yoyotte, 1972; Quentin *et al.*, 1976), so the morphology of the group is better understood.

It appears that the structure of the ovejector is of importance, as this can be either of two distinct types: it may be long (1500-5000  $\mu\text{m}$ ) often with the middle part dilated to form a chamber (e.g. *A. travassosi* Caballero, 1938) or very short (200-600  $\mu\text{m}$ ) and simple (e.g. *A. turgida* Stossich, 1902, in Chabaud and Choquet, 1955). Examination of well described species shows that the character "ovejector long" is found in species with large caudal papillae, and the character "ovejector short" is found in species with small papillae.

This division is borne out by other characters: spicules equal or unequal, with tips wide or fine, presence or absence of deirids, large or small eggs, larvae with long smooth tails (Chabaud *et al.*, 1959) or short spiny tails (Quentin *et al.*, 1976).

The two groups represent Linstow's two genera, *Aprocta* and *Lissonema*, and *Lissonema* is now reinstated. *Aprocta* Linstow, 1883: ovejector short; caudal papillae small; spicules generally subequal with distal extremity finely pointed; deirids absent; eggs small (less than 55  $\mu\text{m}$  long); larvae short (about 200  $\mu\text{m}$ ), with short tail with terminal spines. Type species; *A. cylindrica* Linstow, 1883 (redescribed by Quentin *et al.*, 1976) from Passeriformes.

Other species sufficiently known to be allotted to the genus, with the bird groups in which they occur, are:

- A. angolica* (Vuylsteke, 1953); Passeriformes
- A. caballeri* Ybarra, 1948; Falconiformes and Passeriformes
- A. cercomelae* Sonin, 1966; Passeriformes
- A. colaptidis* Schuurmans-Stekhoven, 1952; Passeriformes
- A. crassa* Railliet et Henry, 1910; Ralliformes
- A. laeviculis* (Sandground, 1933); Passeriformes
- A. matronensis* Railliet et Henry, 1910; Passeriformes
- A. narium* Linstow, 1901; Falconiformes
- A. ophthalmophaga* Stossich, 1902; Falconiformes
- A. orbitalis* Linstow, 1901; Falconiformes
- A. piloscelidis* Schuurmans-Stekhoven, 1952; Charadriiformes

*A. sudarikowi* Sobolev, 1947\*; Charadriiformes et Ralliformes

*A. textori* Vuylsteke, 1953, Passeriformes

*A. turgida* Stossich, 1902 (redescribed by Chabaud et Choquet, 1955); Lariformes.

The species *A. caudata* Mendonca, 1961, of which only the male is known, seems to belong to the Splendidofilariinae.

*Lissonema* Linstow, 1903: ovejector long; at least one large caudal papilla (a single precloacal papilla, 0-4 pairs cloacal papillae); spicules generally equal, with distal extremity wide and blunt; deirids present; eggs large (over 65  $\mu\text{m}$  long), larvae large (over 400  $\mu\text{m}$  long), tail long unarmed. Type species: *L. rotundata* Linstow, 1903 (redescribed by Rasheed, 1960) parasitic in Cuculiformes.

Other species which are sufficiently known to be allotted to the genus, listed with the bird groups in which they occur, are:

- L. brevicaudata* (Chow, 1939) n.comb. (Strigiformes).
- L. calliderma* (Schmidt et Kuntz, 1970) n.comb. (Cuculiformes).
- L. caprimulgi* (Kazubski, 1958) n.comb. (Caprimulgiformes)
- L. cosmetocephala* (Yoyotte, 1972) n.comb. (Cuculiformes)
- L. ghesquieri* (Ezzat et Tadros, 1958) n.comb. (Caprimulgiformes).
- L. golvani* (Diaz-Ungria, 1963) n.comb. (Piciformes)
- L. lepidogrammi* (Tubangui et Masilungan, 1937) n.comb. (Cuculiformes.)
- L. noctuae* (Spaul, 1927) n.comb., redescribed by Chabaud (1951) (Strigiformes).
- L. nyctidromi* (Caballero et Peregrina, 1938) n.comb. (Caprimulgiformes).
- L. obtusa* (Dujardin, 1845) n.comb. (= *Eucampus obtusus* D., 1845, *Eucampus* Chevr., 1833 preoccupied, Coleoptera) (Caprimulgiformes).
- L. papillosa* (Chabaud, Anderson et Brygoo, 1959) n.comb. (Cuculiformes).
- L. proctata* (Lent et Freitas, 1948) n.comb. (Strigiformes).
- L. semenovi* (Skrjabin, 1934) n.comb. (Caprimulgiformes).

\**A. sudarikowi* Sobolev, 1947 (in Sonin, 1966) is hard to classify: according to this author, the spicules are pointed and the eggs small as in *Aprocta* spp., but the caudal papillae are large as in *Lissonema* spp. The ovejector cannot be analysed. However, Sandground (1933) describing analogous material, emphasised the absence of caudal papillae. It seems to us that this species belongs to *Aprocta*.

*L. spasskyi* (Sonin, 1966) n.comb. (Cuculiformes).

*L. travassosi* (Caballero, 1938) n.comb. (Trogoniformes).

The genus *Squamofilaria* Schmerling, 1925, which was reinstated by Anderson and Chabaud (1958) for Aproctinae with a short well sclerotized buccal capsule, is itself heterogenous. In these species, the buccal capsule appears to be a secondary cuticularisation of the wall of the buccal cavity, as in *Lissonema* sp., described below from *Ninox novaeseelandiae*. It is poorly limited externally, as compared with the true buccal capsule seen in *Dipetalonema* spp., *Litomosa* spp., etc. which are descended from forms with a buccal capsule formed from rhabdions.

The species attributed to *Squamofilaria* fall into the same two morphological groups found in *Aprocta* and *Lissonema*, and can be allotted to one or other of these genera as follows:

(a) *Aprocta*

*A. vestibulata* (Johnston et Mawson, 1950) n.comb. (Passeriformes).

*A. pillersi* (Yorke et Maplestone, 1926) n.comb. (Passeriformes).

(b) *Lissonema*

*L. coraciae* (Gmelin, 1790) n.comb. syn. *Squamofilaria coraciae* (Gmelin, 1870). This is the type species of *Squamofilaria* which now falls as a synonym of *Lissonema* (Coraciadiformes).

*L. sicki* (Strachan, 1957) n.comb. (Strigiformes).

*L. nepalensis* (Soota et Chaturverdi, 1970) n.comb. (Coraciadiformes).

*L. ungriai* (Guerrero, 1971)\* n.comb. (Trogoniformes).

It appears that the host ranges for *Lissonema* and *Aprocta* are mutually exclusive.

*Lissonema* spp. are known from the Strigiformes, the Cuculiformes†, the Piciformes (Bucconidae), the Trogoniformes, the Caprimulgiformes and the Coraciadiformes. It is of interest to note that Berlioz, in his Classification of Birds, treated these six groups consecutively (pages 930 to 974, in Grassé, 1950).

*Aprocta* spp. are known from the Lariformes, the Charadriiformes, the Ralliformes (Burrhinidae and

Otididae), the Falconiformes (similarly grouped by Berlioz, p. 884-929), and the vast group of Passeriformes.

The Australian species which have been studied herein are assigned to the two genera *Aprocta* and *Lissonema*.

*Aprocta boulengeri* n.sp.

(Figs. 3, 4)

Material

*Strepera graculina* Shaw (Cracticidae) from Warwick, Qd., from head and nasal cavities, holotype ♀, allotype ♂ (SAM V2098 and V2099) and 1 entire ♀ and 2 broken ♂ (AHC 5839).

Morphology

The morphology is shown in Figures 3 and 4. Unpaired portion of ovejector short and simple.

Dimensions

Holotype ♀: length 50 µm; width 920 µm; cephalic papillae form area 110 µm (laterally) by 70 µm (dorsoventrally); nerve ring and vulva respectively 175 µm and 940 µm from head; oesophagus 1060 µm long, with muscular part 215 µm long; tail length 385 µm, egg 58 µm by 32 µm.

Allotype ♂: length 25.6 mm, diameter 600 µm, cephalic papillae area 110 µm x 55 µm; buccal capsule 20 µm long; nerve ring and excretory pore 205 µm and 305 µm from head; oesophagus 800 µm long with muscular part 190 µm; tail length 320 µm; right and left spicules 400 µm and 440 µm long, 22 µm wide.

Other entire female: body length 37 mm, diameter 910 µm; cephalic papillae on area 88 µm x 57 µm; buccal capsule 38 µm long; nerve ring, excretory pore and vulva respectively 220 µm, 310 µm and 680 µm from head; oesophagus 965 µm long, with muscular part 200 µm; unpaired part of ovejector 600 µm long; tail length 240 µm; larvae 190 and 205 µm long, 11 µm wide.

Discussion

This species is easily differentiated by the following characters: presence of a median subterminal papilla on the male tail, spicules 400-440 µm in length, with tricuspid distal extremities, vulva slightly in front of the oesophago-intestinal junction, anus of the female not terminal. This is proposed as a new species *A. boulengeri*.

*Aprocta bakeri* n.sp.

(Fig. 5)

Material

*Corvus orru* Bonaparte (Corvidae) from Emu Vale, near Warwick, Qd., from nasal cavity: holotype ♂ (SAM V21000).

\* The eggs measure only 45 µm x 25 µm, but their concave form indicates that they are not embryonated.

† Among species from Cuculiformes, those from *Centropus* are unusual in having a long oesophagus (5-7 mm), clearly divided.




Figure 3. *Aprocta boulengeri* n.sp., female, and male allotype. A, anterior end of female, lateral view; B, C, & D, head of female, in apical, lateral and dorsal views; E, F, head of male, ventral and lateral views; G, anterior end of male, lateral view; H, I, tail of female, lateral and ventral views; J, K, L, tail of male, lateral and ventral views; M, left spicule. A, G, H, I, J, L, to scale 200  $\mu\text{m}$ ; B, C, D, E, F, K, to scale 100  $\mu\text{m}$ ; M, to scale 150  $\mu\text{m}$ .




Figure 4. *Aprocta boulengeri* n.sp. (cont.) A, and B, tips of left and right spicules, ventral views; C, larva; D, E, head of larvae, with hooks seen in profile and full face; F, tip of tail of larva. A-E, to scale 10  $\mu\text{m}$ ; F, free-hand.

Description

Morphology is shown in Figure 5. Triangular buccal cavity. Cloaca opening on conspicuous protuberance.

Dimensions

Body length 29 mm, diameter 550  $\mu\text{m}$ ; buccal capsule 40  $\mu\text{m}$  long; nerve ring 140  $\mu\text{m}$  from head; left and right spicules respectively 390  $\mu\text{m}$  and 330  $\mu\text{m}$  long; tail length 260  $\mu\text{m}$ .

Discussion

The bifid extremity of the testis in these specimens resembles that of *Aprocta laevicutis* from *Cyornis banyumas whitei* but is distinguished from this species by the larger body size and by the spicule length (390 and 330  $\mu\text{m}$  instead of 210-170  $\mu\text{m}$ ). It is also distinguished from all other *Aprocta* spp. by a

voluminous protuberance bearing the cloacal opening. It is proposed as a new species, *A. bakeri*, dedicated to our Canadian colleague, Michael Baker.

*Lissonema* sp.

(Fig. 6)

Material

From *Ninox novaeseelandiae* Gmelin (Strigidae) from Berrimah, N.T., 1 ♀ (AHC 6261).

Description

The morphology is shown in Fig. 6. Mouth and buccal capsule narrower dorso-ventrally than laterally; buccal capsule stoutly built, of two parts—anteriorly, a thickening of the cephalic cuticle,




Figure 5. *Aprocta bakeri* n.sp., holotype male. A, anterior end; B, oesophageal region, lateral view; C, D, head in dorso-ventral and lateral views; E, F, tail, in lateral and subventral views; G, H, right and left spicules, ventral views; I, J, distal extremities of same spicules; A, to scale 500  $\mu\text{m}$ ; B, E, F, to scale 200  $\mu\text{m}$ ; C, D, to scale 100  $\mu\text{m}$ ; G, H, to scale 150  $\mu\text{m}$ ; I, J, to scale 10  $\mu\text{m}$ . v, vulva.



posteriorly, a pre-oesophageal thickening, poorly defined externally, and thicker laterally. Base of buccal cavity formed by the three prominent and sclerotised oesophageal lobes. The cuticular 'scales' are irregularly distributed over the body length, but the lateral fields are smooth. Unpaired portion of ovejector long and complex (Fig. 6 J).

#### Measurements

Body length 34 mm, diameter 590  $\mu\text{m}$ ; buccal cavity 22  $\mu\text{m}$  long; nerve ring, excretory pore and deirids at 155  $\mu\text{m}$ , 242  $\mu\text{m}$  and 270  $\mu\text{m}$  from head; oesophagus 1200  $\mu\text{m}$  long, with muscular part 230  $\mu\text{m}$  long; vulva 700  $\mu\text{m}$  from head; unpaired part of ovejector 4660  $\mu\text{m}$  long; tail length 95  $\mu\text{m}$ ; cuticular

scales on body 22  $\mu\text{m}$  long, 13  $\mu\text{m}$  wide, commencing at 1150  $\mu\text{m}$  from head and ending 550  $\mu\text{m}$  from posterior end of body. Eggs 66  $\mu\text{m}$  x 38  $\mu\text{m}$ ; larva 415  $\mu\text{m}$  long, 13.5  $\mu\text{m}$  in diameter.

#### Discussion

Among the species of this genus in which there is a buccal capsule, *L. sicki*, *L. nepalensis*, *L. ungriai* and *L. coraciae*—only the last named agrees with the Australian specimens in having 'scales' on the body, but the shape and arrangement of these was not exactly described. The dimensions of the female are very close, except the length, which is only half that of the Australian specimens. In view of this, and of the different geographical range, no specific identification is made.




Figure 6. *Lissonema* sp., female. A, oesophageal region, lateral view; B, anterior end, median view; C, D, head, lateral and median views; E, cuticular ornamentation in region of mid-body, in submedian view; F, deirid, lateral view; G, cuticular scale in lateral view; H, I, tail, ventral and lateral views; J, ovejector, distal part; K, larva; L, head of larva; A, B, E, H, I, to scale 200  $\mu\text{m}$ ; C, D, F, G, K, to scale 100  $\mu\text{m}$ ; J, to scale, 500  $\mu\text{m}$ ; L, to scale 50  $\mu\text{m}$ .



## II. Diplotriaenoidea—Dicheilonematinae

### Genus *Serratospiculum* Skrjabin

The genus *Serratospiculum* contains (a) species with small spicules (left spicules 600-700  $\mu\text{m}$ , right spicule 300-350  $\mu\text{m}$ ), which may be considered as grouped about *S. guttatum* and (b) species with large spicules (left spicule 1100-1300  $\mu\text{m}$ , right spicule 350-500  $\mu\text{m}$ ) grouped about *S. tendo*.

#### (a) Species with small spicules

*S. guttatum* (Schneider, 1866): the most characteristic feature of this species is the cuticular ornamentation of the female body, in the form of large bosses, well described by Schneider. The species is recorded in Australia from *Falco berigora* by Schneider, 1866, from *F. longipennis* and *F. peregrinus* (Syn. *F. melanogenys*) by Johnston & Mawson, 1941, and in the present paper.

Species grouped with *S. guttatum* may have (i) smooth or (ii) ornamented cuticle.

#### (i) With smooth cuticle:

*S. chungii* Hoeppli & Hsü, 1929, from *Falco* sp., China.

*S. congolensis* Vuylsteke, 1957, from *Butastur rufipennis* (Sund.) from equatorial Africa\*. This species is distinguished from *S. chungii* by the structure of the right spicule: the oblique crests are present on the distal third of the spicule, not on the distal half.

*S. seurati* n.sp. (syn. *Filaria attenuata* Rud., 1819 sensu Seurat, 1915) from various Falconidae, from North Africa, is close to *S. chungii* but can be distinguished by the cephalic structure—the epaulettes concave with regard to the dorso-ventral axis, and not rectilinear (Fig. 18 of Hoeppli & Hsü, 1929), relatively narrower—(30  $\mu\text{m}$  wide by 100  $\mu\text{m}$  long in *S. seurati*, 50  $\mu\text{m}$  wide by 120  $\mu\text{m}$  long in *S. chungii*).

#### (ii) With cuticular ornamentation:

*S. turkestanicum* Skrjabin, 1916 (type species of the genus), from *F. tinnunculus* L. from Central Asia, is well distinguished by the type of cuticular ornamentation, consisting of points curved backwards.

*S. kwangsiensis* Hsü, 1963, from *F. columbarius insignis* (Clark) from China, is close to *S. guttatum* in the structure of the head and the presence of rounded bosses on the cuticle of the female, but these bosses continue to the level of the anus in this

species but only to 1 mm from the posterior extremity in *S. guttatum*; also, the distal half of the spicule is particularly delicate.

The filarial worms described by Li (1933) from *F. columbarius* (syn. *Hypotrionchis aesolon*), from China, as *S. turkestanicum* seems from the character of the female cuticle, to be *S. kwangsiensis*.

#### (b) Species with large spicules

This group contains three species, all without cuticular ornamentation in the female:

*S. tendo* Nitzsch, 1819†, the species is parasitic in *F. peregrinus* in Europe (Nitzsch in Rudolphi, 1819; Dujardin, 1845; Yorke & Maplestone, 1926) and in Australia (material described here and, as pointed out by Sonin (1968), some parts of the material seen by Johnston & Mawson, 1941, and noted as *S. guttatum*). The species is also found in other falcons, and in other parts of the Old World.

*S. thoracis* Tubangui, 1934, from *Falco ernesti* in the Philippines; the dimensions of this species are similar to those of *S. tendo*.

*S. lii* Ezzat & Tadros, 1958, from *F. peregrinus callidus* from Belgian Congo. In this species the measurements resemble those of the two preceding species. The *area rugosa* and the detailed structure of the spicules have not been described, and as pointed out by Sonin (1968), it is possible that these three species are synonymous.

It appears that there are more species of *Serratospiculum* than allowed by Sonin (1968). It also appears that *S. guttatum* is found only in Australia, and that references to its presence in other parts of the world are erroneous.

The Australian material from falcons (Falconidae) permits the redescription of two species, *Serratospiculum guttatum* (Schneider, 1866) and *S. tendo* (Nitzsch, 1819). Comparison of the Australian material with specimens in MNHN has led to the identification of a new species, *S. seurati*.

### *Serratospiculum guttatum* (Schneider, 1866)

(Fig. 7)

#### Material

From *Falco longipennis* Swainson, from Fleurieu Peninsula and Yorke Peninsula, South Australia. 9♂, 5 anterior and 2 posterior ends of ♂, 7♀, 4 anterior and 5 posterior ends ♀ (AHC 6279); 2♂ and 2♀ (MNHN, 193 HD).

From *Falco peregrinus* Tunstall, from Mallala, S.A. 3♂, 1 posterior end ♂, (AHC 6280).

#### Description

The morphology is shown in Fig. 7. The shape of the head is the same in all specimens, . . . broader on lateral axis, epaulettes twice as long as wide (80  $\mu\text{m}$

\* The apical view shown in Fig. 36 appears to be in error; the piece of the head shown is probably seen upside down, from the posterior aspect.

† Railliet & Henry (1916) have resolved clearly the question of the nomenclature of this species.




Figure 7. *Serratospiculum gutatum* A-G, female H-N male, A, anterior end, lateral view; B, cuticular ornamentation in mid-body, lateral view; C, the same, detail of one boss; D, distal end of ovejector; E, F, tail, in lateral and ventral view; G, head, apical view; H, anterior end, ventral view; I, head, median view; J, K, left and right spicules; L, *area rugosa*, ventral view; M, the same, in detail; N, tail, ventral view; A, scale 500  $\mu\text{m}$ ; B, D, H, to scale 200  $\mu\text{m}$ ; C, E, F, to scale 150  $\mu\text{m}$ ; G, I, L, M, N, to scale 75  $\mu\text{m}$ .

by 40  $\mu\text{m}$ ) and in apical view, convex with regard to the dorso-ventral axis. Cuticular ornamentation present only in the female. *Area rugosa* forms two latero-ventral bands of low cuticular protuberances.

#### Dimensions

Three females: length 127, 150 and 140  $\mu\text{m}$ , width 700, 790, and 600  $\mu\text{m}$ ; nerve ring at 175, 180, and 240  $\mu\text{m}$  from head; excretory pore at 205, ?, and 240  $\mu\text{m}$  from head; deirids at 250, 220, and 510  $\mu\text{m}$  from head; muscular oesophagus 420  $\mu\text{m}$  long, and with glandular part 13 mm long (measured on one of the

females dissected); vulva at 740, 600, and 1050  $\mu\text{m}$  from head; length of unpaired part of ovejector 2150  $\mu\text{m}$  (in female dissected); tail 90-110  $\mu\text{m}$  long; embryonated eggs 52 by 30  $\mu\text{m}$ ; cuticular ornamentation starts at 3 mm from the head and ends at 1 mm from the posterior end.

Two males: length 72 and 93 mm, width 480, 560  $\mu\text{m}$ ; nerve ring and excretory pore 150, 230  $\mu\text{m}$  from head (measured on one male); deirids at 280 and 340  $\mu\text{m}$ , 212 and 275  $\mu\text{m}$  from head; muscular oesophagus 320, 365  $\mu\text{m}$  long, length of glandular

oesophagus not determined; tail length 80, 82  $\mu\text{m}$ ; length of left spicules 650, 670  $\mu\text{m}$ , of right spicules 350, 360  $\mu\text{m}$ ; *area rugosa* lying from 5.4 mm to 450  $\mu\text{m}$  from the posterior end (measured on one male).

**Serratospiculum tendo** (Nitzsch, 1819)

(Fig. 8)

**Material**

From *Falco peregrinus* Tunstall, from Port Augusta, S.A. 1 ♀, 1 ♂ (AHC 6273).

This material has been compared with MNHN 510 JJ collected in France from the same host species; this consists of one anterior end and two posterior ends of ♂ and the anterior and posterior ends of 1 ♀.

**Description**

The morphology of the two lots of material is similar, and is shown in Figure 8. The cephalic structure is constant: rounded head, epaulette 3-4 times as long as wide (110  $\mu\text{m}$  by 30  $\mu\text{m}$ ), concave




Figure 8. *Serratospiculum tendo*. A, anterior end of female, lateral view; B, anterior end of male, median view; C, head of female, apical view; D, E, head of male, median and lateral views; F, male, *area rugosa*, ventral view; G, H, female, tail, ventral and lateral views; I, J, tail of male, ventral and lateral views; F and J, Australian material, other figures French material. A, B, G, J, to scale 200  $\mu\text{m}$ ; C, D, E, F, I, to scale 100  $\mu\text{m}$ ; H, to scale 500  $\mu\text{m}$ .

with regard to the dorso-ventral axis, and prolonged by 4 small additional swellings. *Area rugosa* in form of two ventro-lateral bands of small longitudinal cuticular wrinkles. Spicules not drawn, as studied by Bain & Vassiliades (1969).

#### Measurements of Australian specimens

Female: length of body 190  $\mu\text{m}$ , width 825  $\mu\text{m}$ ; nerve ring and deirids 300  $\mu\text{m}$  and 210-220  $\mu\text{m}$  from head; length of oesophagus 19.7 mm, of muscular part 600  $\mu\text{m}$ ; vulva 1650  $\mu\text{m}$  from head; tail 100  $\mu\text{m}$  long.

Male: length of body 148 mm, width 410  $\mu\text{m}$ ; nerve ring and deirids 200  $\mu\text{m}$  and 305-310  $\mu\text{m}$  from head; length of oesophagus 11 mm, of muscular part 600  $\mu\text{m}$ ; length of left spicule 1120  $\mu\text{m}$ , of right 505  $\mu\text{m}$ ; tail 130  $\mu\text{m}$  long; *area rugosa* extends from 4650  $\mu\text{m}$ -600  $\mu\text{m}$  from the caudal extremity.

#### Measurements of specimens from France

In female, vulva 1 mm from the head, tail 140  $\mu\text{m}$  long. In male nerve ring and deirids 165  $\mu\text{m}$  and 230  $\mu\text{m}$  from head; length of left spicule 1150  $\mu\text{m}$ , and 1120  $\mu\text{m}$ , of right spicules 510  $\mu\text{m}$  and 500  $\mu\text{m}$ ; tail 145 and 110  $\mu\text{m}$ .

#### *Serratospiculum seurati* n.sp.

(Fig. 9)

Syn. *Filaria attenuatum* Rudolphi, 1819 *sensu* Seurat, 1915.

#### Material

Specimens studied by Seurat, 1915, and determined as '*Filaria attenuata* Rud., 1819 (= *F. guttatum* Schneider, 1866)'. This material, now in the MNHN, Paris, was taken from *Falco biarmicus erlangeri* Kleinschm at Biskra, North Africa; it comprises 2 $\delta$  of which one is in two pieces, 1




Figure 9. *Serratospiculum seurati* n.sp. A, anterior end of female, lateral view; B, head of holotype female, lateral view; C, head of female, apical view; D, E, tail of holotype female, ventral and lateral views; F, tail of male, subventral view; G, H, left and right spicules, lateral view. A, E, to scale 200  $\mu\text{m}$ ; B, C, D, F, H, to scale 100  $\mu\text{m}$ ; G, to scale 50  $\mu\text{m}$ .

anterior end ♂, 2 ♀ and 2 posterior ends ♀. 1 entire ♀ and 1 entire ♂ were chosen as holotype and allotype respectively (MNHN 110 E1).

(2) specimens collected by Dollfus (MNHN 114 A) in 1914 from *Falco* sp., from Rabat, Morocco, labelled *S. guttatum* (Schneider, 1866); consisting of numerous pieces including 2 anterior and 4 posterior ends of ♀, 3 anterior and 1 posterior ends ♂.

#### Morphology

The general morphology, well described by Seurat, is complemented by Figure 8. The structure of the head, in both lots of material, is similar. The position of the vulva is somewhat variable (450-1000 µm from head). An *area rugosa* appears to be absent. The right spicule bears low crests, obliquely placed.

#### Measurements

Holotype ♀: length of body 135 mm, width 770 µm; vulva 470 µm from head; tail length 40 µm. Specimen not sufficiently clear for other measurements.

Allotype ♂: length of body 42 mm, width 480 µm; nerve ring and deirids 130 µm and 230 µm from head; length of oesophagus 10 mm, of its muscular part 270 µm; left spicule and right spicules respectively 775 µm and 300 µm; tail length 110 µm.

From MNHN 114 A (2 pieces): length unknown; width 500 µm; nerve ring 170 µm from head; muscular oesophagus 310 µm; left and right spicules respectively 650 µm and 350 µm, tail length 110 µm.

#### Diplotriaeonoidea: Diplotriaeonidae

##### *Diplotriaeona spratti* n.sp.

(Fig. 10)

#### Material

From *Oreoica gutturalis* Vigors & Horsfield (Muscicapidae) from the Petermann Ranges, N.T. Holotype ♀ and allotype ♂ (SAM, V2101 and V2102 respectively); 6 ♀, 9 ♂, and numerous pieces of ends of ♀ and 1 posterior end ♂ (AHC 6272); 1 ♀ and 2 ♂ paratypes (MNHN, Paris, 204 HD).

#### Description

Morphological features are shown in Figure 10.

#### Measurements

Holotype ♀: length of body 68 mm, width 700 µm; nerve ring 240 µm from head; tridents 150 µm

and 172 µm long; length of oesophagus 3720 µm, of muscular part 370 µm; vulva 500 µm from head, unpaired portion of ovejector 3300 µm long; tail length 110 µm; eggs, non-fertilized, 32 µm by 18 µm.

Paratype ♀s: body length 66-71 mm; in ♀ length 66 mm, width 500 µm; nerve ring and vulva 290 µm and 425 µm from head; trident 135 µm; oesophagus 4500 µm; eggs 45 µm by 35 µm.

Allotype ♂: body length 32 mm, width 400 µm; nerve ring 200 µm from head; muscular oesophagus 3850 µm long; tail 50 µm; left and right spicules respectively 4500 and 600 µm long.

Paratype ♂s: length of body 32-34 mm; of oesophagus 2600-3275 µm; left spicule 4200-6300 µm, right spicule 510-600 µm; in one male 32 mm long; body width 455 µm, nerve ring 230 µm from head; tridents 125 µm; length of oesophagus 3400 µm, of tail 55 µm; left spicule 4200 µm long, with calamus 650 µm, right spicule 550 µm.

#### Discussion

This species can be compared with the others of the genus in which the left spicule is longer than 2000 µm. These are: *D. bhamaensis* (Parona, 1889)\* redescribed by Anderson (1959) from Sturnidae and Turdidae in Asia, has the trident not pointed but concave at the apex, a much longer oesophagus (8-12 mm instead of 2.6-6.3 mm), left spicule not exceeding 2600 µm, and the right spicule spiralled about a strongly curved axis, whereas in the Australian specimens the axis is almost straight.

In *D. ecaudata* (Oerley, 1882), from Sturnidae in Africa the trident is concave at the tip, the left spicule is not longer than 2800 µm, and the axis of the right spicule has three strong bends.

*D. tricuspis* (Fedtschenko, 1874) sensu Singh, 1949, is, as pointed out by Anderson (1959), a mistaken determination. It is clear from Singh's paper that though the female is a *Diplotriaeona* sp., the male (Fig. 27) is a *Hamatospiculum* sp. The Australian species is proposed as *D. spratti* n.sp., named after our colleague David Spratt. The species is characterised by the pointed apex of the trident, the long left spicule (over 4000 µm) and the right spicule spiralled around an almost straight axis.

\* *B. buckleyi* Fotedar & Kaw, 1965, and *D. mirzapurensis* Soota & Chaturvedi, 1967, both parasites of *Acridotheres tristis*, have no morphological difference from *D. bhamaensis*, and are regarded as synonyms of this species.




Figure 10. *Diplotriaena spratti* n.sp. A, anterior end of holotype female, lateral view; B, head of holotype female, median view; C, junction of oesophagus and intestine, paratype female; D, E, tail of paratype female, lateral and ventral views; F, trident of holotype female; G, to Q, male: G, anterior end, ventral view; H, head, apical view; I, J, tridents of two specimens; K, posterior end, lateral view; L, M, the two extremities of the left spicule; N, right spicule and part of the left spicule; O, distal end of right spicule, ventral view; P, Q, tail in lateral and ventral views. A, D, to scale 500  $\mu\text{m}$ ; B, to scale 150  $\mu\text{m}$ ; C, E, G, L, N, to scale 200  $\mu\text{m}$ ; F, I, J, M, O, P, Q, to scale 100  $\mu\text{m}$ .

### *Diplotriaena beveridgei* n.sp.

(Fig. 11)

#### Material

From *Corvus orru* Bonaparte (Corvidae) from Emu Vale, near Warwick Qd: Holotype ♀ (SAM V 2103) and posterior end (10 mm) of allotype ♂ (SAM V 2104); 2 ♀, 2 anterior and 3 posterior ends ♀ (AHC 6276).

#### Description

Morphological features are shown in Figure 11.

#### Measurements

Holotype ♀: body length 102 mm, width 1400  $\mu\text{m}$ , nerve ring 450  $\mu\text{m}$  from head; trident 285  $\mu\text{m}$  long; oesophagus 9500  $\mu\text{m}$  long, its muscular part 575  $\mu\text{m}$ ; vulva 630  $\mu\text{m}$  from head; eggs 50 x 35  $\mu\text{m}$ ; tail 200  $\mu\text{m}$  long.

Allotype ♂: the piece is 1.11 mm wide; tail 10  $\mu\text{m}$  long; left spicule 1355  $\mu\text{m}$  long (calamus 250  $\mu\text{m}$ ); right 1075  $\mu\text{m}$ .

Paratypes: one anterior end of ♀, 1400  $\mu\text{m}$  wide, with nerve ring 390  $\mu\text{m}$  from head, tridents 250  $\mu\text{m}$




Figure 11. *Diplotriaeana beveridgei* n.sp. A-H, female; I-L, male. A, anterior end, lateral view; B, oesophago-intestinal junction; C, head, apical view; D, head, lateral view; E, F, trident and its apex; G, H, tail, ventral and lateral views; I, posterior end, ventral view; J, K, right and left spicules; L, tail, ventral view. A, B, G, H, to scale 500  $\mu$ m; C, to scale 150  $\mu$ m; D, E, J, K, to scale 200  $\mu$ m; F, L, to scale 100  $\mu$ m; I, to scale 1000  $\mu$ m.

and 270  $\mu$ m; oesophagus 8950  $\mu$ m with muscular part 580  $\mu$ m; vulva 650  $\mu$ m from head, unpaired ovejector 2900  $\mu$ m. One posterior end with tail 200  $\mu$ m. Length of entire female 81 mm.

Discussion

The specimens differ from the other two *Diplotriaeana* spp. from Australian Corvidae as follows:

In *D. alpha* Johnston & Mawson, 1940, the trident is smaller (120  $\mu$ m instead of 250-285  $\mu$ m) and oesophagus twice as long.

In *D. beta* Johnston & Mawson, 1940, considered by Anderson (1959) as a synonym of *D. flabellata* (Linstow, 1888) the oesophagus is very long, but the tridents shorter (190-200  $\mu$ m), more massive, and with rounded apex, and the right spicule is a different shape.



The right spicule is bent in an 'L' similar to that of *D. delirae* Pinto & Noronha, 1970, from Tyrannidae in Brazil, however our material is distinguished from this species by the longer oesophagus, and the longer right spicule (1075  $\mu\text{m}$  instead of 610  $\mu\text{m}$ ).

This material is therefore proposed as *D. beveridgei* n.sp., named after the collector, Dr Ian Beveridge. It is characterised by the large tridents, the long oesophagus (about 9 mm) and the 'L' shape of the right spicule.

***Diplotriaeana smithi* n.sp.**

(Fig. 12)

**Material**

5 collections from *Acanthogenys rufogularis* Gould: From The Bunkers, S. Aust.: 4 birds, as follows:

1. holotype ♀ (SAM V2105), allotype ♂ (SAM V2106)  
2 ♀, 1 ♂, pieces of ♀ and ♂ (AHC 6278)  
1 ♀, 1 ♂, posterior end 1 ♂ (MNHN, Paris 208 HD)
2. 1 ♀, 1 ♂ pieces of ♀ (AHC 6270)
3. 1 ♀, pieces of ♀ and ♂ (AHC 6268)
4. 4 ♀, 3 ♂, pieces of ♀ (AHC 6267)

From Wubin, W. Aust.: 6 ♀, 3 ♂ (AHC 6264); 3 ♀, 2 ♂ (MNHN, 208 HD).

**Description**

Morphological details are shown in Figure 12.

**Dimensions**

Holotype ♀: body length 50 mm, width 680  $\mu\text{m}$ ; trident length 220  $\mu\text{m}$ ; nerve ring and vulva 350  $\mu\text{m}$  and 660  $\mu\text{m}$  from head; length of oesophagus 4000  $\mu\text{m}$ , of tail 45  $\mu\text{m}$ , of unpaired part of ovejector 2100  $\mu\text{m}$ .

Allotype ♂: length of body 22 mm, width 530  $\mu\text{m}$ ; trident length 190  $\mu\text{m}$ ; nerve ring and excretory pore 310  $\mu\text{m}$  and 350  $\mu\text{m}$  from head; length of oesophagus 2850  $\mu\text{m}$ , of tail 68  $\mu\text{m}$ , of left spicule 800  $\mu\text{m}$ , of right spicule 650  $\mu\text{m}$ .

Paratype 3 ♀s: Length of body 35, 22, 39 mm; width 650, 610, 520  $\mu\text{m}$ ; tridents 205, 195, 190  $\mu\text{m}$ ; nerve ring 300, 290, 330  $\mu\text{m}$  from head; length of oesophagus 2400, 2100  $\mu\text{m}$ , damaged in 3rd; vulva 650, 730, 500  $\mu\text{m}$  from head; tail length 40, 55, 55  $\mu\text{m}$ .

Paratype ♂: body length 22 mm, width 520  $\mu\text{m}$ , trident length 165  $\mu\text{m}$ ; nerve ring 280  $\mu\text{m}$  from head; oesophagus 3400  $\mu\text{m}$  long (muscular part 340  $\mu\text{m}$ ); length of tail 80  $\mu\text{m}$ ; of left spicule 785  $\mu\text{m}$ , of right spicule 500  $\mu\text{m}$ .

Variations observed among all the specimens: tridents 165-190  $\mu\text{m}$  in males, 190-220  $\mu\text{m}$  in females; vulva 480-730  $\mu\text{m}$  from head; oesophagus 2000-3600  $\mu\text{m}$  long in ♀; left spicule 730-850  $\mu\text{m}$ , right spicule 500-680  $\mu\text{m}$ . One male from AHC 6278 is teratological, with left and right spicules respectively 260  $\mu\text{m}$  and 310  $\mu\text{m}$ .

**Discussion**

*D. zeta* Johnston & Mawson, 1940, redescribed by Anderson (1959) was also taken from *Acanthogenys rufogularis*; the only female known is of similar dimensions to the present specimens, except that the trident is smaller (150  $\mu\text{m}$  instead of 190-205  $\mu\text{m}$ ). Moreover, there are two grooves on the head of tridents in *D. zeta* not seen on any of the new specimens. These two differences, and the absence of a male of *D. zeta*, prevent the identification of the new material as *D. zeta*.

Five other species must be considered:

Two species have similar dimensions and spicule shape: *D. clelandi* (Johnston, 1912) from Cracticidae in Australia (male only known) and *D. kennedyi* Grewal, 1965, from Pycnonotidae from India. These two species are of doubtful status, as their morphology is imperfectly known, and it is impossible to compare our specimens with them.

The three other species in which the trident and spicules are similar to our specimens are distinguished as follows: in *D. pungens* (Schneider, 1866) from Turdididae in Asia, the oesophagus is very long (14500  $\mu\text{m}$  in the ♀) and three pairs of papillae lie distinctly in front of the cloaca. In *D. flabellata* (Linstow, 1888), from Paradiseidae in the Aru Islands, the trident is more stoutly built, the body is particularly wide, the glandular oesophagus very long (6500-7500  $\mu\text{m}$  in the female)—the left spicule strongly curved and the male tail rounded.

In *D. urocissoides* Anderson, 1959, from Corvidae in India, the head structure is characteristic ('buccal cavity surrounded by a clearly defined cuticular thickening often forming a plug,' Anderson, 1959) and the number of papillae is small.

It seems that the material described here is new, and the name *D. smithi* is proposed for it, in honour of our colleague I. Humphrey Smith. It is characterised by the large trident with blunt tip and without grooves, the oesophagus not very thick and not particularly long or short (2000-4000  $\mu\text{m}$ ), the spicules of medium length (730-850  $\mu\text{m}$  and 500-600  $\mu\text{m}$ ) the tail of male rectangular in ventral view, with precloacal papillae few in number and situated close to the cloaca.




Figure 12. *Diplotrriaena smithi* n.sp. A, anterior end of female holotype, lateral view; B, head of female paratype, apical view; C, D, same female, apex of tridents; E, F, tail of same female in lateral and ventral views; G, anterior end of male allotype, lateral view; H, same male, oesophago-intestinal junction; I, head of same male, lateral view; J, trident of male paratype; K, tail of male allotype, ventral view; M, N, male allotype, right and left spicules, ventral views; O, P, two other views of same right spicule; A, E, G, M, N, O, P, to scale 200  $\mu\text{m}$ ; B, H, F, J, K, L, to scale 100  $\mu\text{m}$ ; C, D, to scale 50  $\mu\text{m}$ ; I, to scale 150  $\mu\text{m}$ .

***Diplotrriaena delta* Johnston & Mawson, 1940**  
(Fig. 13 A-O)

Material

From *Amytornis goyderi* (Gould) (Maluridae), from the Simpson Desert, S. Aust.: 10 ♀, 2 ♂ (AHC 6274); 2 ♀ and 1 ♂ (MNHN 207 HD).

Description

Additional morphological details are given in Figure 13 A-O.

Dimensions

2 ♀: body length 70, 72 mm, width 460, 400  $\mu\text{m}$ ;

tridents 115-120  $\mu\text{m}$  and 100  $\mu\text{m}$ ; nerve ring 180, 190  $\mu\text{m}$  from head; oesophagus length 1850, 2110  $\mu\text{m}$ , its muscular part 330, 160  $\mu\text{m}$ ; vulva 525, 400  $\mu\text{m}$  from head; tail length 70, 32  $\mu\text{m}$ ; eggs 46 x 29  $\mu\text{m}$ . The extremes of the oesophagus are 1750  $\mu\text{m}$  and 2400  $\mu\text{m}$ , in females measuring respectively 62 mm and 70 mm.

2 ♂: body length 30, 28 mm, width 400, 380  $\mu\text{m}$ ; tridents 110, 120  $\mu\text{m}$ ; nerve ring 170, 160  $\mu\text{m}$  from head; left spicule 950  $\mu\text{m}$  long, (calamus 240  $\mu\text{m}$ ), and 1050  $\mu\text{m}$ ; right spicule 720, 630  $\mu\text{m}$  long; tail 50, 60  $\mu\text{m}$ .

## Discussion

These specimens are identified as *D. delta*, described from *Malurus lamberti* (Maluridae) in Australia. This species, placed by Anderson (1959) as a synonym of *D. tridens* (Molin, 1858), was reinstated by Ogden (1965), who had specimens from *M. lamberti* from various parts of Australia.

*D. delta* is recognisable by the simultaneous presence of the following characters (taken from the

new material and from Ogden): large body size (60-127 mm in female); tridents small (78-122  $\mu\text{m}$ ) with three small digitations at the tip; oesophagus of average size (1750-3900  $\mu\text{m}$  in female, 1560-2600  $\mu\text{m}$  in male), with glandular part thin or thick; left spicule 950-1140  $\mu\text{m}$ , right spicule 620-730  $\mu\text{m}$ , with triple twist; tail of male with two lateral swellings.




Figure 13. A-O, *Diplotrinaea delta*, A-E, female: A, anterior end, subventral view; B, head, en face; C, D, trident and its apex; E, tail, ventral view; F-O, male: F, anterior end; G, H, trident and its apex; I, J, two views of right spicule; K, L, right and left spicules of another male, lateral view; M, N, tips of right and left spicules, ventral views; O, tail, ventral view.

P-R, *D. falconis*. P, tail of female, ventral view; Q, profile of cuticle at mid-body; R, tail of male, ventral view.

A, F, Q, to scale 500  $\mu\text{m}$ ; I, J, K, L, to scale 200  $\mu\text{m}$ ; P, to scale 125  $\mu\text{m}$ ; B, E, G, M, N, O, R, to scale 100  $\mu\text{m}$ ; H, to scale 50  $\mu\text{m}$ .

**Diplotriaeana falconis** (Connal, 1912)

(Fig. 13 P-R)

## Material

From *Falco berigora* Vigors & Horsfield, from Yorke Peninsula, S. Aust.: 3♀, 1♂ (AHC 6269).

## Description

Additional morphological details are given in Figure 13 P-R.

## Measurements

2♀: body length 70, 72 mm; width 590, 660 µm; trident length 135, 135 µm; nerve ring 230, 230 µm from head; vulva 430, 660 µm from head; oesophagus in second female 2550 µm long, with muscular part 400 µm; tail 22, 28 µm; eggs 65 x 30 µm.

♂: body length 45 mm, width 550 µm; trident length 135 µm, oesophagus length 2700 µm, with muscular part 450 µm; tail 32 µm long; left spicule broken, 1025 µm long; right 1100 µm long (850 µm if measured in a straight line).

## Discussion

The measurements, general morphology, and the characteristic anatomy of the left spicule agree with those of *D. falconis*, redescribed by Anderson (1959), from Falconidae in Africa. In the Australian specimens, there are ten pairs of papillae in the male, the tail of the female bears papillae, and the cuticle of the female is marked with regular and distinct undulations.

## ACKNOWLEDGEMENTS

We are grateful to all those who collected the birds hosts or the parasites which enabled this work to be undertaken . . . these are Dr Ian Beveridge, Mrs Joan Paton, Dr Dom Serventy, Mr Shane Parker, and officials of the South Australian Museum and the Northern Territory Administration (Animal Industry and Agriculture Branch).

## APPENDIX

Australian Species Described in this Paper, Arranged under their Hosts

## Corvidae

*Corvus orru* Bonaparte  
*Aprocta bakeri* n.sp.  
*Diplotriaeana beveridgei* n.sp.

## Cracticidae

*Strepera graculina* (Shaw)  
*Aprocta boulengeri* n.sp.

## Muscipapidae

*Petroica multicolor* (Gmelin)  
*Pseudaprocta copemani* n.sp.  
*Pachycephala pectoralis* (Latham)  
*Pseudaprocta copemani* n.sp.  
*Oreoica gutturalis* (Vigors and Horsfield)  
*Diplotriaeana spratti* n.sp.

## Maluridae

*Amytornis gwyderi* (Gould)  
*Diplotriaeana delta* Johnston and Mawson

## Meliphagidae

*Acanthogenys rufogularis* Gould  
*Diplotriaeana smithi* n.sp.

## Falconidae

*Falco berigora* (Vigors and Horsfield)  
*Diplotriaeana falconis* (Connal)  
*Serratospiculum guttaum* (Schneider)  
*Falco peregrinus* Tunstall  
*S. tendo* (Nitzsch)  
*Falco longipennis* Swainson  
*Serratospiculum guttaum* (Schneider)

## Strigidae

*Ninox novaeseelandiae* (Gmelin)  
*Lissonema* sp.

## REFERENCES

- ALI, S. M. 1956. Studies on the nematode parasites of fishes and birds found in Hyberabad State. *Indian J. Helminth.*, **8**, 1-83.
- ANDERSON, R. C. 1957. The life cycles of Dipetalonematid Nematodes (Filarioidea, Dipetalonematidae): the problem of their evolution. *J. Helminth.*, **31**, 203-224.
- ANDERSON, R. C. and CHABAUD, A. G. 1958. Taxonomie de la Filaire *Squamofilaria sicki* (Strachan, 1957) n. comb. et place du genre *Squamofilaria* Schermerling, 1925 dans la sousfamille Aproctinae. *Annls Parasit. hum. comp.*, **33**, 254-266.
- ANDERSON, R. C. 1959. Preliminary revision of the genus *Diplotriaeana* Henry et Ozoux, 1909 (Diplotriaeinae: Diplotriaeinae). *Parassitologia*, **1**, 195-314.
- ANDERSON, R. C. and BAIN, O. 1976. CIH Keys to the Nematode Parasites of Vertebrates. No. 3. Keys to genera of the Order Spirurida, Part 3. Diplotriaeinoidea, Aproctoidea and Filarioidea. Ed. by R. C. Anderson, A. G. Chabaud and S. Willmott.
- BAIN, O., and VASSILIADIS, G. 1969. Cycle évolutif d'un Dicheil-onematinae, *Serratospiculum tendo*, Filaire parasite du Faucon. *Annls Parasit. hum. comp.*, **44**, 595-604.
- BERTHOZ, J. 1950. Systématique. In Grassé P. P., *Traité de Zoologie, Oiseaux*, **15**, 1164 pp., Paris, Masson Edit.
- BOULENGER, M. A. 1928. Report on a collection of parasitic Nematodes, mainly from Egypt, part V. Filarioidea. *Parasitology*, **20**, 32-55.
- CABALLERO, E. y C. 1938. Contribucion al conocimiento de los Nematodos de las aves de México. V. Libro Jubilar Prof. Travassos. Rio de Janeiro, 91-97.
- CHABAUD, A. G. 1951. Observation sur *Aprocta noctuae* Spaul, 1927 (Nematoda—Aproctidae). *Arch. Inst. Pasteur Maroc*, **4**, cah. 3; 236-243.
- CHABAUD, A. G. 1974. C.I.H. Keys to the Nematode parasites of Vertebrates; No. 1. Keys to Subclasses, Orders and Superfamilies. Ed. by R. C. Anderson, A. G. Chabaud, S. Willmott.
- CHABAUD, A. G., ANDERSON, A. C. and BRYGOO, E. R. 1959. Sept Filaires d'Oiseaux malgaches. *Annls Parasit. hum. comp.*, **39**, 69-94.
- CHABAUD, A. G., BRYGOO, E. R. and RICHARD, J. 1964. Filaires d'Oiseaux malgaches (Deuxième note). *Ann. Parasit. hum. comp.*, **39**, 69-94.
- CHABAUD, A. G. and CHOQUET, M. T. 1955. Helminthes de la région de Banyuls. II. Deux Filaires parasites d'Oiseaux. *Vie et Milieu*, **6**, 93-100.
- DIAZ-UNGRÍA, C. 1963. Nematodes parasites, nouveaux ou intéressants, du Venezuela. *Annls Parasitol. hum. comp.*, **38**, 893-913.
- DUJARDIN, F. 1845. Histoire naturelle des Helminthes ou Vers intestinaux. Paris, 654 pp.
- EZZAT, M. A. E. et TADROS, G. 1958. Contribution to the helminth fauna of Belgian Congo birds. *Annls Mus. r. Congo Belge*, sér. 8, **69**, 1-81.
- FOTEDAR, D. N. and KAW, L. 1965. New Diplotriaeinid nematode from the body cavity of *Acridotheres tristis* (Linnaeus). *Kashmir Sci.*, **2**, 125-131.
- HÖEPLI, R. and HSU, H. F. 1929. Helminthologische Beiträge aus Fukien und Chekiang. *Arch. Schiffsh. Tropenhyg.*, **33**, 1-43.
- HSU, W. N. 1963. (Studies on some parasitic Nematodes of order Spirurida Chitwood, 1933). *Acta zool. sin.*, **15**, 544-552.

- JOHNSTON, T. H. 1912. Notes on some Entozoa. *Proc. Roy. Soc. Qland*, **24**, 63-91.
- JOHNSTON, T. H. and MAWSON, P. M. 1940. Some filarial parasites of Australian birds. *Trans. R. Soc. South Aust.*, **64**, 355-361.
- JOHNSTON, T. H. and MAWSON, P. M. 1941. Some Nematodes from Australian birds of prey. *Trans. R. Soc. South Aust.*, **65**, 30-35.
- LI, H. C. 1933. Report on a collection of parasitic nematodes, mainly from North China. 1. Filarioidea. *Parasitology*, **25**, 192-224.
- LINSTOW, O. 1883. Nematoden, Trematoden und Acanthocephalen, gesammelt von Prof. Fedtschenko in Turkestan. *Arch. Naturg., Berlin*, **49**, 274-314.
- LINSTOW, O. 1903. Parasiten, meistens Helminthen, aus Siam. *Arch. Mikr. Anat.*, **62**, 108-121.
- OGDEN, C. G. 1966. Some parasitic Nematodes from Australian birds. *Annl's Mag. Nat. Hist.*, ser. 13, **9**, 505-518.
- PINTO, R. M. and NORONHA, D. 1970. Sobre uma nova especie do género *Diplotrriaena* Railliet et Henry, 1909 (Nematoda, Filarioidea) *Diplotrriaena delirae* sp. n. *Atas Soc. Biol. Rio de Janeiro*, **14**, 55-57.
- QUENTIN, J. C., TRONCY, P. M. and BARRE, N. 1976. *Aprocta cylindrica* Linstow, 1883, Filaire ovipare parasite d'Oiseaux Ploceides au Tchad, Morphogénèse larvaire du Nématode. *Annl's Parasitol. hum. comp.*, **51**, 83-93.
- RAILLIET, A. and HENRY, A. 1916. Les Filaires des Rapaces (Falconiformes et Strigiformes). *Bull. Soc. Path. Exot.*, **9**, 364-368.
- RASHEED, S. 1960. The Nematode parasites of the birds of Hyderabad (India). *Biologia, Lahore*, **6**, 116 pp.
- RUDOLPHI, C. A. 1819. *Entozoorum synopsis cui accedunt mantissa duplex et indices locupletissimi*. Berolini, 811 pp.
- SANDGROUND, J. H. 1933. Report on the nematode parasites collected by the Kelley-Roosevelts Expedition to Indo-China with descriptions of several new species. *Z. ParasitKde*, **5**, 542-583.
- SCHIKHOBALOVA, N. P. 1930. Sur une nouvelle filaire d'Oiseaux *Pseudaprocta gubernacularia* n.g., n.sp. *Annl's Parasit. hum. comp.*, **8**, 624-627.
- SCHMIDT, G. D. and KUNTZ, R. E. 1970. Nematodes parasites of Oceanica. VIII. Filariids of birds, with a new genus and four new species. *Parasitology*, **60**, 313-326.
- SCHNEIDER, A. 1866. Monographie der Nematoden, Berlin, 357 pp.
- SEURAT, L. G. 1915. Expédition de M. M. Walter Rotschild, E. Hartert et C. Hilgert dans le Sud-Algérien (Mars-Mai 1914). Nématodes parasites. *Novitates Zoologicae*, **22**, 1-25.
- SINGH, S. N. 1949. Studies on the helminth parasites of birds in Hyderabad State. Nematoda IV. *J. Helminth.*, **23**, 39-56.
- SKRJABIN, K. I. 1915. Nématodes des Oiseaux du Turkestan russe. *Ann. Mus. Zool. Acad. St. Petersbourg*, **20**, 531-557.
- SONIN, M. D. 1966. (Principles of nematology, edited by K. I. Skrjabin. Vol. 17. Filariata of animals and man and the diseases caused by them. Part 1, Aproctoidea). Moscow: Izdatelstvo "Nauka", 360 pp. (In Russian).
- SONIN, M. D. 1968. *Osnovi Nematodologii*, **21**, Diplotrriaenoidea, Acad. Sci. Edit., Moscow, pp. 388.
- SOOTA, T. D. and CHATURVEDI, Y. 1967. On two new Nematode species of the genus *Diplotrriaena* Railliet et Henry, 1909, from Nepal and India. *J. Zool. Soc. India*, **19**, 133-136.
- TUBANGUI, M. A. 1934. Nematodes in the collection of the Philippine Bureau of Science, II: Filarioidea. *Philippine J. Sc.*, **55**, 115-123.
- VUYLSTEKE, C. 1957. Nématodes parasites d'Oiseaux. Exploration du Parc National de la Garamba. Mission H. de Saeger. Bruxelles, **8**, 3-20.
- YORKE, W. and MAPLESTONE, P. A. 1926. The Nematode parasites of Vertebrates. London, J & A Churchill, 536 pp.
- YOYOTTE, E. V. 1972. Etude de huit Nématodes parasites de Vertébrés du Venezuela et de la Colombie. *Bull. Mus. nat. Hist. nat.*, 3e sér., **41**, Zool. 35, 477-498.